

The topology of the category of open and closed strings

Nils A. Baas Ralph L. Cohen ^{*} Antonio Ramírez

February 1, 2008

Dedicated to Sam Gitler on the occasion of his 70th birthday.

Abstract

In this paper we study the topology of the cobordism category $\mathcal{S}^{oc}$ of open and closed strings. This is a 2-category in which the objects are compact one-manifolds whose boundary components are labeled by an indexing set (the set of “ D -branes”), the 1-morphisms are cobordisms of manifolds with boundary, and the 2-morphisms are diffeomorphisms of the surface cobordisms. Our methods and techniques are direct generalizations of those used by U. Tillmann in her study of the category of closed strings. We input the striking theorem of Madsen and Weiss regarding the topology of the stable mapping class group to identify the homotopy type of the geometric realization $|\mathcal{S}^{oc}|$ as an infinite loop space.

Introduction

As described by Segal [7], in a conformal field theory one associates to a closed 1-dimensional manifold S a Hilbert space $\mathcal{H}_S$, and to a 2-dimensional cobordism Σ between closed 1-manifolds S_1 and S_2 , an operator $\varphi_\Sigma : \mathcal{H}_{S_1} \rightarrow \mathcal{H}_{S_2}$. This operator depends on the conformal class of a metric on the surface Σ . Such a field theory can be viewed as a representation of the category $\mathcal{S}$ whose objects are closed 1-manifolds, and where the space of morphisms from S_1 to S_2 is the moduli space of conformal classes of metrics (i.e Riemann surfaces) on cobordisms from S_1 to S_2 .

In [8] U. Tillmann described $\mathcal{S}$ as a 2-category, whose objects are nonnegative integers, corresponding to the number of components in the closed 1-manifold, the 1-morphisms between m and n are cobordisms between a disjoint union of m -circles and a disjoint union of n -circles, and whose 2-morphisms are diffeomorphisms of the cobordisms that are fixed on the boundary. She constructed $\mathcal{S}$ as a symmetric monoidal 2-category, and with a clever argument using the Harer stability theorem, she proved that there is a homotopy equivalence

$$\mathbb{Z} \times B\Gamma_\infty^+ \simeq \Omega|\mathcal{S}|.$$

^{*}The second author was partially supported by a grant from the NSF

Here $|\mathcal{S}|$ is the bisimplicial geometric realization (classifying space) of the category $\mathcal{S}$, and $\Omega|\mathcal{S}|$ is its based loop space. $B\Gamma_\infty^+$ is Quillen's plus construction on the classifying space of the stable mapping class group,

$$B\Gamma_\infty = \varinjlim_g B\Gamma_{g,1}$$

where $\Gamma_{g,n}$ is the mapping class group of surfaces of genus g with n boundary components. Since $\mathcal{S}$ is a symmetric monoidal category this result established $B\Gamma_\infty^+$ as an infinite loop space.

This result indicated the important role of infinite loop space theory in studying the stable mapping class group, and it eventually led to Madsen and Weiss's proof of a generalization of a conjecture of Mumford [3], saying that there is an equivalence of infinite loop spaces

$$B\Gamma_\infty^+ \simeq \Omega^\infty \mathbb{CP}_{-1}^\infty.$$

Here $\Omega^\infty \mathbb{CP}_{-1}^\infty$ is the zero space of the stunted projective spectrum $\mathbb{CP}_{-1}^\infty$, which is defined to be the Thom spectrum of minus the canonical line bundle $L \rightarrow \mathbb{CP}^\infty$, $\mathbb{CP}_{-1}^\infty = (\mathbb{CP}^\infty)^{-L}$.

The purpose of this paper is to generalize Tillmann's results to the category $\mathcal{S}^{\text{oc}}$ of "open and closed strings." In this category the objects are compact 1-manifolds which may have boundary. In other words, the objects are disjoint unions of circles and closed intervals. The need to study such a category arises from open-closed string theory, in the presence of " D -branes" (see for example, [6]). If the background space of the string theory is a manifold M , the D -branes are submanifolds $B_i \subset M$ that define the boundary conditions of the strings. In other words, one is considering spaces of closed and "open" strings moving in M , where the open strings are maps from the interval $\gamma : [0, 1] \rightarrow M$, with boundary conditions $\gamma(0) \in B_0$, and $\gamma(1) \in B_1$, where B_0 and B_1 are D -branes. In practice, D -branes are often equipped with bundles and connections, but we will not be concerned with this additional structure in this paper.

In order to encode the additional information the presence of D -branes imposes, we consider an abstract finite set $\mathcal{B}$ which is meant to index the set of D -Branes in the background manifold M , and we define the objects of $\mathcal{S}_{\mathcal{B}}^{\text{oc}}$ to be compact 1-manifolds (unions of circles and intervals) where each interval has its boundary points labeled by an element of $\mathcal{B}$. These labels are meant to represent the boundary values of maps from these intervals.

The 1-morphisms in this category will be "open-closed cobordisms." These are cobordisms between the compact manifolds with boundary that respect the boundaries of the compact one-manifolds (in the sense that the cobordisms restrict to give ordinary cobordisms of the boundaries of the 1-manifolds), as well as the labeling by elements of $\mathcal{B}$. In other words, the boundary of an open-closed cobordism is partitioned into three parts:

1. the "incoming part," consisting of a number of circles and intervals, whose boundaries are labeled by elements of $\mathcal{B}$,
2. the "outgoing part" of the same type, and,

3. the “free boundary,” which is itself a cobordism between the boundaries of the incoming and outgoing parts. Each connected component of the free boundary is labeled by an element of $\mathcal{B}$, which respects the labeling of the boundaries of the incoming and outgoing parts.

Finally, the 2-morphisms are diffeomorphisms of the cobordisms that respect all of this structure.

To state our main result, we introduce some notation. Let $F_{g,n}$ be an oriented surface of genus g with n boundary components. Let $\mathbf{m}$ be a fixed, finite set of points in $F_{g,n}$. The set $\mathbf{m}$ is partitioned into subsets indexed by $\mathcal{B}$,

$$\mathbf{m} = \bigcup_{b \in \mathcal{B}} m_b.$$

We think of the subset m_b as those points in the set $\mathbf{m}$ that are “colored” by the element $b \in \mathcal{B}$. Let $\text{Diff}(F_{g,n}, \mathbf{m}, \partial)$ be the group of orientation-preserving diffeomorphisms of $F_{g,n}$ that fix the boundary pointwise and preserve each of the sets m_b for $b \in \mathcal{B}$. Let $\Gamma_{g,n}^{\mathbf{m}} = \pi_0 \text{Diff}(F_{g,n}, \mathbf{m}, \partial)$ denote the corresponding mapping class group.

Let $B\Gamma_{\infty,n}^{\mathcal{B}}$ be the colimit of the classifying spaces, $\varinjlim_{g,\mathbf{m}} B\Gamma_{g,n}^{\mathbf{m}}$. Here we are taking the limit over the genus g and the sets $\mathbf{m}$, which are partially ordered by inclusion. In this limit, the cardinality of each subset m_b tends to infinity. The following is the main theorem of this paper.

Theorem 1. *1. The category $\mathcal{S}_{\mathcal{B}}^{oc}$ of open and closed strings with set of D -branes $\mathcal{B}$ can be given the structure of a symmetric monoidal 2-category, so its geometric realization $|\mathcal{S}_{\mathcal{B}}^{oc}|$ is an infinite loop space.*

2. For any n there is a homology equivalence

$$\mathbb{Z} \times \mathbb{Z}^{|\mathcal{B}|} \times B\Gamma_{\infty,n}^{\mathcal{B}} \xrightarrow{\cong_{H*}} \Omega|\mathcal{S}_{\mathcal{B}}^{oc}|.$$

Here $|\mathcal{B}|$ is the cardinality of the set $\mathcal{B}$.

3. There is a homotopy equivalence of infinite loop spaces,

$$\Omega|\mathcal{S}_{\mathcal{B}}^{oc}| \simeq \Omega^{\infty} \mathbb{CP}_{-1}^{\infty} \times \prod_{b \in \mathcal{B}} Q(\mathbb{CP}_+^{\infty})$$

where, as usual, X_+ denotes X with a disjoint basepoint and $Q(Y) = \varinjlim_k \Omega^k \Sigma^k(Y)$.

The paper is organized as follows. In Section 1 we define the notion of open-closed cobordism carefully and define the category $\mathcal{S}^{oc}$ corresponding to the case when there is only one D -brane. That is, the set $\mathcal{B}$ consists of one element. This simplifying assumption will allow us not to be concerned with the labeling of the boundaries of the 1-manifolds. In this section we also show that the category $\mathcal{S}^{oc}$ has a symmetric monoidal structure. As the reader will see, our constructions are direct generalizations of those of Tillmann [8], [9]. However the objects in our category will have to be stratified in a rather delicate way, in order to keep track of the boundary configurations in the cobordisms. In Section 2 we adapt Tillmann’s argument involving Harer stability to prove

part 2 of Theorem 1 when there is only one D -brane. In this setting, part 3 of Theorem 1 will follow immediately from part 2 using the Madsen-Weiss theorem [3], and a result of Bödighheimer and Tillmann [1] which we review. In Section 3 we describe the necessary modifications to prove Theorem 1 for a general set $\mathcal{B}$ of D -branes.

1 Open-closed cobordisms and the category $\mathcal{S}^{\text{oc}}$.

We begin by defining the notion of an open-closed cobordism.

Definition 1. A smooth (two-dimensional) open-closed cobordism $\Sigma = (\Sigma, \partial^\pm \Sigma, \partial_{\text{f}} \Sigma)$ is a smooth oriented surface Σ , together with a choice of subsets $\partial^- \Sigma$, $\partial^+ \Sigma$, $\partial_{\text{f}} \Sigma$ of its boundary $\partial \Sigma$, such that

1. $\partial \Sigma = \partial^- \Sigma \cup \partial^+ \Sigma \cup \partial_{\text{f}} \Sigma$,
2. each of $\partial^- \Sigma, \partial^+ \Sigma, \partial_{\text{f}} \Sigma$ is an embedded one-dimensional manifold with boundary (to which we assign the orientation induced from $\partial \Sigma$), and,
3. $\partial_{\text{f}} \Sigma$ is a cobordism from $\partial(\partial^- \Sigma)$ to $\partial(\partial^+ \Sigma)$.

The sets $\partial^- \Sigma$, $\partial^+ \Sigma$, $\partial_{\text{f}} \Sigma$ are called the incoming, outgoing, and free boundaries, respectively. We will write $\partial^\pm \Sigma$ for $\partial^- \Sigma \cup \partial^+ \Sigma$.

Given two such cobordisms Σ_1 and Σ_2 , we say that they are isomorphic if there exists an orientation-preserving diffeomorphism $\varphi : \Sigma_1 \rightarrow \Sigma_2$ which restricts to diffeomorphisms $\partial^- \Sigma_1 \rightarrow \partial^- \Sigma_2$, $\partial^+ \Sigma_1 \rightarrow \partial^+ \Sigma_2$, $\partial_{\text{f}} \Sigma_1 \rightarrow \partial_{\text{f}} \Sigma_2$. Denote the space of such isomorphisms by $\text{Diff}_{\text{oc}}^+(\Sigma_1, \Sigma_2)$. If we also have given diffeomorphisms $\partial^- \Sigma_1 \cong \partial^- \Sigma_2$ and $\partial^+ \Sigma_1 \cong \partial^+ \Sigma_2$, we define $\text{Diff}_{\text{oc}}^+(\Sigma_1, \Sigma_2; \partial^\pm)$ to be the set of elements of $\text{Diff}_{\text{oc}}^+(\Sigma_1, \Sigma_2)$ which restrict to the fixed diffeomorphisms on $\partial^\pm \Sigma_1 = \partial^\pm \Sigma_2$.

We now go about defining the category of open and closed strings. Actually there will be several versions of this category which will have different applications.

The objects in such a category are essentially diffeomorphism classes compact 1-manifolds, which may or may not have boundary. Such a manifold is diffeomorphic to a disjoint union of circles and intervals. However, as we will see below, the order of these manifolds, and even various groupings of the intervals, will be important. So we will define a “general object” to be a sequence of diffeomorphism classes of circles and intervals, which we enumerate by sequences of zeros and ones (standing for the circles and intervals, respectively).

Definition 2. A general object is a sequence $\vec{n} = (n_1, \dots, n_k)$, where $n_i \in \{0, 1\}$. The length of the sequence k can be an arbitrary nonnegative integer. Each such tuple $\vec{n}$ stands for a one-dimensional submanifold $C_{\vec{n}} = \bigcup_{i=1}^k I(i, n_i)$ of $\mathbb{R}^2$, where we define $I(i, 0)$ to be the circle of radius $1/4$ centered at $(i, 0)$ and $I(i, 1) = [i - 1/4, i + 1/4] \times \{0\}$. We will use the notation $\ell(\vec{n}) = k$ for the length of a tuple $\vec{n}$.

A general morphism between $\vec{n}$ and $\vec{m}$ is an open-closed cobordism Σ smoothly embedded in $\mathbb{R}^3$, with $\partial^- \Sigma = C_{\vec{n}} \times \{0\}$ and $\partial^+ \Sigma = C_{\vec{m}} \times \{t\}$ for some $t > 0$. In addition, we require that:

1. $\partial^\pm \Sigma$ has a collar, in such a way that for some $\mu > 0$ we have $C_{\vec{n}} \times [0, \mu), C_{\vec{m}} \times (t - \mu, t] \subset \Sigma$.
2. The orthogonal projection in the t -direction is a Morse function on the interior of Σ .
3. The same projection is a Morse function when restricted to $\partial_f \Sigma$.




Figure 1: An open-closed cobordism from $\vec{n} = (1, 1, 1, 0, 1)$ to $\vec{m} = (0, 0, 1, 1, 0)$

The first condition in the definition of general morphism allows us to define compositions in the following way. Given $\Sigma_1 \subset \mathbb{R}^2 \times [0, t_1]$ a general morphism from $\vec{n}$ to $\vec{m}$, and $\Sigma_2 \subset \mathbb{R}^2 \times [0, t_2]$ a general morphism from $\vec{m}$ to $\vec{k}$, define their composition $\Sigma_2 \circ \Sigma_1 \subseteq \mathbb{R}^2 \times [0, t_1 + t_2]$ as the union of Σ_1 with the result of translating Σ_2 by t_1 in the t -direction. This clearly yields an associative composition.

The categories of open and closed strings we will consider will actually be 2-categories. We therefore need the following definition.

Definition 3. Given Σ_1 and Σ_2 general morphisms from $\vec{n}$ to $\vec{m}$, we define the space of general 2-morphisms, $2\text{-Mor}_{gen}(\Sigma_1, \Sigma_2) = \emptyset$ unless Σ_1 and Σ_2 are isomorphic, in which case we define $2\text{-Mor}_{gen}(\Sigma_1, \Sigma_2)$ to be the set of $\varphi \in \text{Diff}_{oc}^+(\Sigma_1, \Sigma_2; \partial^\pm)$ which fix some collar of $\partial^\pm \Sigma_1 \cong \partial^\pm \Sigma_2$, where we identify collars of the target 1-manifolds of Σ_1 and Σ_2 by a suitable linear scaling in the z direction.

Notice that by composition of diffeomorphisms, for each pair of general objects $\vec{n}$ and $\vec{m}$, we have a category $\mathcal{S}_{gen}^{oc}(\vec{n}, \vec{m})$ of general morphisms and 2-morphisms. Furthermore, composition of morphisms defines functors

$$\mathcal{S}_{gen}^{oc}(\vec{n}, \vec{m}) \times \mathcal{S}_{gen}^{oc}(\vec{m}, \vec{k}) \longrightarrow \mathcal{S}_{gen}^{oc}(\vec{n}, \vec{k})$$

which together define a 2-category $\mathcal{S}_{gen}^{oc}$ with objects, 1-morphisms, and 2-morphisms given by the general objects, general morphisms, and general 2-morphisms defined above.

We now define certain partitions associated to our general objects. We do this by considering a connected open-closed cobordism (general morphism) Σ from $\vec{n}$ to (0) . Here the general object (0) is a sequence of length one and corresponds to a single circle.

Let $\partial_I \Sigma$ be the set of those connected components of $\partial^\pm \Sigma$ which are intervals. In the case we are considering all of $\partial_I \Sigma$ lies in the incoming part, $\partial^- \Sigma$, and are in bijection with the “1” entries in the sequence $\vec{n}$. Notice that the cobordism Σ defines a partition of $\partial_I \Sigma$ into subsets consisting of those intervals that lie in the same path component of the boundary, $\partial \Sigma$.

We describe this combinatorially as follows.

Definition 4. Let k be the length of a general object $\vec{n}$. Let $I(\vec{n}) = (i_1, \dots, i_{\alpha(\vec{n})})$ be the ordered subset of $\{1, \dots, k\}$ defined by $j \in I(\vec{n})$ if and only if $n_j = 1$. So the length $\alpha(\vec{n})$ of $I(\vec{n})$ is the number of ones in the sequence $\vec{n}$.

Consider the permutation $\sigma_\Sigma(\vec{n})$ of $I(\vec{n})$ defined as follows. Let $j \in I(\vec{n})$. By definition this represents an interval $I_j \subset \partial_I \Sigma$. I_j lies in a connected component of the topological boundary $\partial \Sigma$. Define $\sigma_\Sigma(j) = \ell$, where I_ℓ is the interval lying next to I_j in this component, in the ordering induced by the orientation of Σ . We call this permutation an open boundary permutation of $\vec{n}$, or more precisely, of the one-manifold $C_{\vec{n}} = \partial^+ \Sigma$. We view this permutation σ_Σ as an element of the symmetric group $S_{\alpha(\vec{n})}$.




Figure 2: $\vec{n} = (0, 1, 1, 1)$, $\sigma = (2, 3)(4) \in S_{\alpha(\vec{n})}$

Notice that different open-closed cobordisms Σ from $\vec{n}$ to (0) can define the same permutation $\sigma \in S_{\alpha(\vec{n})}$, and that every element of the symmetric group can arise in this way.

We now define how open boundary permutations on $\vec{n}$ may be pulled back by a general morphism $\Sigma_{\vec{m}, \vec{n}}$ from $\vec{m}$ to $\vec{n}$.

Definition 5. Let $\Sigma_{\vec{m}, \vec{n}}$ be a general morphism from $\vec{m}$ to $\vec{n}$. Let $\sigma \in S_{\alpha(\vec{n})}$ be the open boundary permutation of $\vec{n}$ and choose a connected general morphism $\Sigma_{\vec{n}, (0)}$ from $\vec{n}$ to (0) inducing σ . Define the pullback permutation $\Sigma_{\vec{m}, \vec{n}}^*(\sigma) \in S_{\alpha(\vec{m})}$ to be the open boundary permutation of $\vec{m}$ induced by the glued cobordism $\Sigma_{\vec{m}, \vec{n}} \circ \Sigma_{\vec{n}, (0)}$ from $\vec{m}$ to (0) .

We leave it as an easy exercise for the reader to check that this notion of pullback is well-defined.

Next, we define a 2-category $\mathcal{S}_D^{\text{oc}}$ as a refinement of $\mathcal{S}_{\text{gen}}^{\text{oc}}$ that incorporates these open boundary permutations.

Definition 6. Define the 2-category $\mathcal{S}_D^{\text{oc}}$ to have objects consisting of pairs $(\vec{n}, \sigma)$, where $\vec{n}$ is a general object and $\sigma \in S_{\alpha(\vec{n})}$. Define the 1-morphisms, $\mathcal{S}_D^{\text{oc}}((\vec{n}, \sigma), (\vec{m}, \tau))$ to be those general morphisms Σ from $\vec{n}$ to $\vec{m}$ such that $\Sigma^*(\sigma) = \tau$. The 2-morphisms from Σ_1 to Σ_2 are the general 2-morphisms, i.e., diffeomorphisms from Σ_1 to Σ_2 as in definition 3.

Define the category $\mathcal{S}_\Gamma^{\text{oc}}$ as the quotient $\mathcal{S}_D^{\text{oc}}/\equiv$, where $\equiv$ is the equivalence relation that identifies pairs of 2-morphisms in the same path component in the corresponding $\text{Diff}_{\text{oc}}^+(\Sigma_1, \Sigma_2; \partial^\pm)$. This category has the same objects and morphisms as $\mathcal{S}_D^{\text{oc}}$, with each space $\text{Diff}_{\text{oc}}^+(\Sigma_1, \Sigma_2; \partial^\pm)$ of 2-morphisms replaced by its set of connected components, the mapping class groups which we denote by $\Gamma(\Sigma_1, \Sigma_2; \partial^\pm)$. We refer to [9] for a treatment of the relevant quotient construction on 2-categories. The following result is immediate.

Proposition 2. The canonical map $\pi_0 : \mathcal{S}_D^{\text{oc}} \rightarrow \mathcal{S}_\Gamma^{\text{oc}}$ is a 2-functor between 2-categories.

Now we describe a pairing on $\mathcal{S}_D^{\text{oc}}$ which is not well-defined, nor associative, nor functorial. However, this pairing will descend to a symmetric monoidal structure on a subquotient of $\mathcal{S}_\Gamma^{\text{oc}}$.

Given $\vec{n}_1, \vec{n}_2 \in \text{Ob}(\mathcal{S}_D^{\text{oc}})$, define $\vec{n}_1 \otimes \vec{n}_2$ to be the juxtaposition of the two tuples with $\vec{n}_1$ followed by $\vec{n}_2$. Given $\Sigma_1 \in \mathcal{S}_D^{\text{oc}}(\vec{n}_1, \vec{m}_1)$, $\Sigma_2 \in \mathcal{S}_D^{\text{oc}}(\vec{n}_2, \vec{m}_2)$, we construct $\Sigma_1 \otimes \Sigma_2 \in \text{Mor } \mathcal{S}_D^{\text{oc}}(\vec{n}_1 \otimes \vec{n}_2, \vec{m}_1 \otimes \vec{m}_2)$ by taking the union of Σ_1 and Σ_2 after they have been modified by suitable diffeomorphisms of $\mathbb{R}^3$ to make them be disjoint and have the same height. More precisely, construct $\Sigma_1 \otimes \Sigma_2$ as follows. Suppose that $\Sigma_i \subset \mathbb{R}^2 \times [0, t_i]$. Choose a collar-preserving diffeomorphism $r : [0, t_2] \rightarrow [0, t_1]$; in this way the diffeomorphism $R(x, y, z) := (x, y, r(z))$ takes Σ_2 to an open-closed cobordism $R(\Sigma_2) \in \mathcal{S}_D^{\text{oc}}(\vec{n}_2, \vec{m}_2)$ which lies in $\mathbb{R}^2 \times [0, t_1]$. Next, choose a smooth orientation-preserving function $s : [0, t_1] \rightarrow \mathbb{R}_{\geq 0}$ which equals $1 + \ell(n_1)$ in a neighborhood of 0 and $1 + \ell(m_1)$ in a neighborhood of t_1 , in such a way that the map $S(x, y, z) := (x + s(z), y, z)$ takes $R(\Sigma_2)$ into an smooth open-closed cobordism $\tilde{\Sigma}_2$ with collar which is disjoint from Σ_1 . Then, define $\Sigma_1 \otimes \Sigma_2 := \Sigma_1 \cup \tilde{\Sigma}_2$. As in the closed case [8] this yields a well-defined map of 2-morphisms which descends to a map of 2-morphisms in $\mathcal{S}_\Gamma^{\text{oc}}$.

To make this pairing into a strictly associative bifunctor, we will make an identification among surfaces. As described for the closed case [9], we will identify two surfaces in $\mathcal{S}_D^{\text{oc}}(\vec{n}, \vec{m})$ if they are

related by a sequence of moves of the same types as those described in Section 3 of [9]. We describe these moves next.

Let $\Sigma \subseteq \mathbb{R}^2 \times [0, t]$ be a smooth open-closed cobordism, and pick n planes $P_k = \mathbb{R}^2 \times \{t_k\}$ where $0 < t_1 < \dots < t_n < t$, in such a way that each of the sets $\Sigma_0, \dots, \Sigma_n$ into which the planes divide Σ is itself a smooth open-closed cobordism between 1-manifolds $\Sigma \cap P_k$ and $\Sigma \cap P_{k+1}$. Here, $P_0 := \mathbb{R}^2 \times \{0\}$, $P_{n+1} := \mathbb{R}^2 \times \{t\}$. Let the connected components of Σ_k be $\Sigma_{k,i}$, $i = 1, \dots, j_k$. Allow each $\Sigma_{k,i}$ to be independently transformed by a sequence of operations of the following two types, as long as the resulting images $\tilde{\Sigma}_{k,i}$ are pairwise disjoint:

- *R-move (Rescaling the height)*: This is an operation of the form $R_{k,i}(x, y, z) = (x, y, r_{k,i}(z))$ on each connected component $\Sigma_{k,i}$, where $r_{k,i} : [t_k, t_{k+1}] \rightarrow [t_k, \tilde{t}_{k,i}]$ is a diffeomorphism with derivative one on a neighborhood of $\partial[t_k, t_{k+1}]$. This rescales the height of all the connected components of Σ_k to make it lie in $\mathbb{R}^2 \times [t_k, \tilde{t}_{k+1}]$; the slices $\Sigma_{k+1}, \dots, \Sigma_n$ are understood to be translated by $\tilde{t}_{k+1} - t_{k+1}$ in the z direction.
- *S-move (Shifting parallel to the xy -plane)*: This is an operation of the form $S_{k,i}(x, y, z) = (x, y, z) + (s_{k,i}^1(z), s_{k,i}^2(z), 0)$, where the $s_{k,i}^j : [t_k, t_{k+1}] \rightarrow \mathbb{R}^2$ are smooth maps which vanish on a neighborhood of $\partial[t_k, t_{k+1}]$. In particular, this move allows passing different connected components of Σ across each other.

Define a 2-category $\mathcal{S}_D^{\text{oc},b}$ with the same objects as $\mathcal{S}_D^{\text{oc}}$, by letting $\mathcal{S}_D^{\text{oc},b}((\vec{m}, \sigma), (\vec{n}, \tau))$ be the full subcategory of $\mathcal{S}_D^{\text{oc}}((\vec{m}, \sigma), (\vec{n}, \tau))$ consisting of those open-closed cobordisms Σ such that no connected component of Σ is an open-closed cobordism to the empty 1-manifold. Define the 2-category $\mathcal{S}_\Gamma^{\text{oc},b}$ analogously, and define $[\mathcal{S}_D^{\text{oc},b}]$, $[\mathcal{S}_\Gamma^{\text{oc},b}]$ to be the quotient categories under the R and S moves.

Theorem 3. *The monoidal pairing above descends to $[\mathcal{S}_\Gamma^{\text{oc},b}]$ and makes it a strict symmetric monoidal 2-category.*

We omit the proof of this theorem, which is a formal argument using the following lemma, proceeding exactly as the proof of Theorem 3.3 in [9].

Lemma 4. *Let $\Sigma \in \mathcal{S}_D^{\text{oc}}((\vec{m}, \sigma), (\vec{n}, \tau))$, and let $E(\Sigma) \subset \text{Diff}_{\text{oc}}^+(\Sigma, \partial^\pm)$ be the subgroup of automorphisms which are obtained as the composition of a sequence of R - and S -moves. Then, $E(\Sigma)$ is contained in the connected component of the identity.*

Proof. The argument is the same as the one in [9] for the closed case. One argues, from the form of the allowable moves, that for Σ connected, any $\varphi \in E(\Sigma)$ must fix each critical point of the t -axis projection, and hence that φ fixes pointwise the subset S of the union of the critical level sets which consists of those connected components containing a critical point. Since the complement of S in Σ is a union of cylinders over closed 1-manifolds with boundary, the result follows because the R - and S -moves cannot twist these cylinders. For Σ not connected but lying in $\mathcal{S}_D^{\text{oc},b}$, the argument

goes through because the condition that no connected component of Σ is a cobordism to the empty 1-manifold guarantees that S -moves cannot swap two connected components of Σ . $\square$

As in [9], given a 2-category $\mathcal{C}$, we denote by $\mathcal{BC}$ the category enriched over topological spaces in which $\text{Ob}(\mathcal{BC}) = \text{Ob}(\mathcal{C})$ and $\mathcal{BC}(x, y)$ equals the realization of the category $\mathcal{C}(x, y)$. We will use $|\mathcal{C}|$ to denote the bisimplicial realization $B\mathcal{BC}$ of the nerve of $\mathcal{BC}$.

For the remainder of the paper, we ease the notation and let the category of open and closed strings, $\mathcal{S}^{oc}$, be the the category $\mathcal{S}_\Gamma^{oc, b}$ above. The following is an immediate consequence of theorem 3 using Segal's well known relationship between symmetric monoidal structures on a category and infinite loop structures on its classifying space [5].

Theorem 5. *The classifying space of the category of open and closed strings, $|\mathcal{S}^{oc}|$, is an infinite loop space.*

2 The homology type of the category $\mathcal{S}^{oc}$

As in the introduction, let $F_{g,n}^m$ denote a fixed oriented surface of genus g , with n boundary components, and m marked points. Let $\text{Diff}(F_{g,n}^{(m)}, \partial)$ be the group of orientation-preserving diffeomorphisms that fix the boundary pointwise, and the marked points only setwise. That is, the diffeomorphisms may permute the marked points. Let $\Gamma_{g,n}^{(m)} = \pi_0(\text{Diff}(F_{g,n}^{(m)}, \partial))$ be the corresponding mapping class group. We consider the colimits of the classifying spaces as g and m increase,

$$B\Gamma_{\infty,n}^{(\infty)} = \varinjlim_{g,m \rightarrow \infty} B\Gamma_{g,n}^{(m)}.$$

The main goal of this section is to prove the following two results.

Theorem 6. *For each n there is a homology equivalence,*

$$\alpha : \mathbb{Z} \times \mathbb{Z} \times B\Gamma_{\infty,n}^{(\infty)} \xrightarrow{\simeq_{H*}} \Omega B\mathcal{S}^{oc}.$$

Theorem 7. *For each n there is a homotopy equivalence*

$$\mathbb{Z} \times \mathbb{Z} \times (B\Gamma_{\infty,n}^{(\infty)})^+ \simeq \Omega^\infty(\mathbb{CP}_{-1}^\infty) \times Q(\mathbb{CP}_+^\infty)$$

where the superscript “+” refers to the Quillen plus construction with respect to the maximal perfect subgroup of the fundamental group.

As we will see below, theorem 7 follows quickly from a splitting theorem for stable mapping class groups due to Bödigheimer and Tillmann [1], and Madsen and Weiss's proof of the stable Mumford conjecture. So most of this section will be devoted to the proof of Theorem 6.

Proof of Theorem 6. Consider the morphism category, $\mathcal{S}_D^{\text{oc}}((\vec{n}, \sigma), (0))$. Remember that the 1-manifold $C_{(0)}$ is a single circle, so that there are no boundary permutations for (0) . Notice that on the classifying space level we have

$$B\mathcal{S}_D^{\text{oc}}((\vec{n}, \sigma), (0)) \simeq \coprod_{\Sigma} B\text{Diff}_{\text{oc}}^+(\Sigma; \partial^{\pm})$$

where Σ ranges over the isomorphism types of open-closed cobordisms from $C_{\vec{n}}$ to $C_{(0)}$, that induce the boundary permutation $\sigma \in S_{\alpha(\vec{n})}$. In the absence of open strings, the isomorphism type of a connected cobordism from n circles to 1 circle is completely determined by its genus. In our present setting, there is another invariant of connected open-closed cobordisms in addition to genus. This is the number w of “windows” of an open-closed cobordism Σ , which is the number of connected components of the free boundary which are homeomorphic to circles. See Figure 3.




Figure 3: An open-closed cobordism with window number $w = 3$.

We remark that if we had not fixed a boundary permutation $\sigma \in S_{\alpha(\vec{n})}$, then the isomorphism type of a connected open-closed cobordism Σ from $\vec{n}$ to (0) would be determined by *three* invariants: its genus, the number of windows, and the boundary permutation $\sigma(\Sigma) \in S_{\alpha(\vec{n})}$.

Let $\Sigma_{(\vec{m}, \tau), (\vec{n}, \sigma)}$ be an open-closed cobordism representing a 1-morphism in $\mathcal{S}_D^{\text{oc}}$. Let $\varphi \in \text{Diff}_{\text{oc}}^+(\Sigma_{(\vec{m}, \tau), (\vec{n}, \sigma)}, \partial^{\pm})$. The connected components of the topological boundary of $\Sigma_{(\vec{m}, \tau), (\vec{n}, \sigma)}$, and the action of φ on them, are given as follows:

1. There are the circle components of the incoming and outgoing boundary. By definition, these are fixed pointwise by φ .
2. There are w windows, i.e., circle components of the free boundary. These are only fixed by φ as a set. It follows that φ is isotopic to a diffeomorphism of the corresponding surface with punctures replacing the “windows”. By a “puncture” we mean a point that has been removed

from the surface, where in the case of a “window”, an open disc has been removed from the surface. Such a diffeomorphism may permute these punctures.

3. Call the components of the topological boundary $\partial\Sigma_{(\vec{m},\tau),(\vec{n},\sigma)}$ that contain at least one incoming or outgoing interval “mixed” boundary components. Since φ fixes each of the incoming and outgoing boundaries pointwise, it follows that φ restricts to a self-diffeomorphism on each of the mixed components. Moreover, since each of these diffeomorphisms fixes a whole interval, φ is isotopic to a diffeomorphism that fixes each of the mixed components pointwise.

From this description, we have that that

$$B\pi_0 \text{Diff}_{\text{oc}}^+(\Sigma_{(\vec{m},\tau),(\vec{n},\sigma)}, \partial^\pm) \simeq B\Gamma_{g,c(\Sigma)}^w,$$

where $c(\Sigma)$ is the number of boundary components of $\partial\Sigma$ of the first or second type, and w is the number of windows, or punctures. In particular for an open-closed cobordism from $(\vec{m},\tau)$ to (0) , the number $c(\Sigma)$ is the number of zeros in the sequence $\vec{m}$ plus the number of disjoint cycles in $\tau \in S_{\alpha(\vec{m})}$, plus one. Since, in this case this number only depends on the object $(\vec{m},\tau)$, we call it $c((\vec{m},\tau))$.

Thus for an object $(\vec{m},\tau)$ of the open-closed string category $\mathcal{S}^{\text{oc}}$, we have proven the following.

Lemma 8. *Let $\mathcal{BS}^{\text{oc}}((\vec{m},\tau), (0))$ be the classifying space of the category of morphisms from the object $(\vec{m},\tau)$ in $\mathcal{S}^{\text{oc}}$ to the object (0) . Then*

$$\mathcal{BS}^{\text{oc}}((\vec{m},\tau), (0)) \simeq \coprod_{w,g \geq 0} B\Gamma_{g,c(\vec{m},\tau)}^w.$$

Fix an open-closed cobordism T in $\mathcal{S}^{\text{oc}}((0), (0))$ with one window and genus one. See Figure 4.




Figure 4: The surface T

By gluing T on the outgoing boundary and extending isomorphisms by the identity, we obtain a functor from $\mathcal{S}^{\text{oc}}((\vec{n},\sigma), (0))$ to itself, and hence a map

$$t : \mathcal{BS}^{\text{oc}}((\vec{n},\sigma), (0)) \longrightarrow \mathcal{BS}^{\text{oc}}((\vec{n},\sigma), (0)). \quad (1)$$

Let $\mathcal{BS}_\infty^{\text{oc}}((\vec{n},\sigma))$ be the homotopy colimit of the system

$$\mathcal{BS}^{\text{oc}}((\vec{n},\sigma), (0)) \xrightarrow{t} \mathcal{BS}^{\text{oc}}((\vec{n},\sigma), (0)) \xrightarrow{t} \dots$$

Since the maps in this colimit increase both the genus and the number of windows, Lemma 1 implies the following.

Lemma 9.

$$\mathcal{BS}_\infty^{\text{oc}}((\vec{n}, \sigma)) \simeq \mathbb{Z} \times \mathbb{Z} \times B\Gamma_{\infty, c(\vec{n}, \sigma)}^{(\infty)}.$$

Notice that in the statement of this lemma, the two copies of $\mathbb{Z}$ reflect the genus and the number of windows.

This definition of extends to a contravariant functor $\mathcal{BS}_\infty^{\text{oc}}$ from $\mathcal{BS}^{\text{oc}}$ to spaces, since, clearly, a morphism ψ from $(\vec{n}, \sigma)$ to $(\vec{m}, \tau)$ defines a pull back map

$$\psi^* : \mathcal{BS}_\infty^{\text{oc}}((\vec{m}, \tau)) \rightarrow \mathcal{BS}_\infty^{\text{oc}}((\vec{n}, \sigma)),$$

which in view of Lemma 9 is homotopic to the gluing map

$$\psi^* : \mathbb{Z} \times \mathbb{Z} \times B\Gamma_{\infty, c(\vec{m}, \tau)}^{(\infty)} \rightarrow \mathbb{Z} \times \mathbb{Z} \times B\Gamma_{\infty, c(\vec{n}, \sigma)}^{(\infty)}.$$

But Harer's stability theorem, as adapted by B\"odigheimer and Tillmann [1] to the setting of infinitely many punctures, implies that each such gluing map is a homology isomorphism. By the generalized group completion theorem ([4], [8]) this implies the following.

Lemma 10. *Let $\text{hocolim} \mathcal{BS}_\infty^{\text{oc}}$ be the homotopy colimit of the functor $\mathcal{BS}_\infty^{\text{oc}}$. Then the natural projection map*

$$p : \text{hocolim} \mathcal{BS}_\infty^{\text{oc}} \rightarrow B\mathcal{S}^{\text{oc}}$$

is a homology fibration, with fiber homology equivalent to $\mathbb{Z} \times \mathbb{Z} \times B\Gamma_{\infty, n}^{(\infty)}$ for any positive integer n .

Notice that $\text{hocolim} \mathcal{BS}_\infty^{\text{oc}}$ is the homotopy colimit under the geometric realization of the maps induced by the operation t of the homotopy colimit of the contravariant functor $\mathcal{BS}_0^{\text{oc}}$ which assigns to an object $(\vec{m}, \tau)$, the space of morphisms, $\mathcal{BS}^{\text{oc}}((\vec{m}, \tau), (0))$. But the homotopy colimit of this functor is contractible since it is the geometric realization of a category with a terminal object (the Quillen over-category $\mathcal{S}^{\text{oc}}/(0)$). Thus $\text{hocolim} \mathcal{BS}_\infty^{\text{oc}}$ is the homotopy colimit of contractible spaces, and hence is itself contractible. (See [8] for this type of argument.)

Thus the homotopy fiber of the map $p : \text{hocolim} \mathcal{BS}_\infty^{\text{oc}} \rightarrow B\mathcal{S}^{\text{oc}}$ is homotopy equivalent to the loop space, $\Omega B\mathcal{S}^{\text{oc}}$. Theorem 6 then follows from Lemma 10. $\square$

We now address Theorem 7. The basis for this theorem is the following result of B\"odigheimer and Tillmann ([1], Theorem 3.1), which makes a clever use of the Harer stability theorem [2].

Theorem 11. *There is a homotopy equivalence,*

$$(B\Gamma_{\infty, n}^{(k)})^+ \simeq (B\Gamma_\infty)^+ \times B(S_k \wr S^1),$$

where S_k is the symmetric group on k letters and $S_k \wr S^1$ is the corresponding wreath product.

We remark that in this theorem $(B\Gamma_\infty)^+$ reflects $(B\Gamma_{\infty,n})^+$ for any n , since as argued in [1], an easy consequence of Harer's stability theorem is that the homotopy type of the plus construction of the classifying space of the stable mapping class group does not depend on how many (fixed) boundary components there are.

For the sake of completeness, we give a sketch of the proof of this theorem. We refer the reader to [1] for details. The idea is to consider the central extension

$$\mathbb{Z}^n \rightarrow \Gamma_{\infty,n} \rightarrow \Gamma_\infty^n$$

which gives rise to a map $\theta : \Gamma_\infty^n \rightarrow (GL^+(2, \mathbb{R}))^n = B(\mathbb{Z}^n) \simeq (S^1)^n$. Here GL^+ stands for matrices having positive determinant. The map θ can be viewed as taking a representative diffeomorphism of a surface with n marked points, $\varphi : F \xrightarrow{\cong} F$, and considering the derivatives, $D\varphi_{x_i} : T_{x_i}F \xrightarrow{\cong} T_{x_i}F$. By fixing framings at the marked points, these can be viewed as elements of $GL^+(2, \mathbb{R})$.

In the mapping class group $\Gamma_\infty^{(n)}$, the representative diffeomorphisms can permute the marked points among themselves, and therefore one has a map

$$\theta_n : \Gamma_\infty^{(n)} \rightarrow S_n \wr GL^+(2, \mathbb{R})$$

which yields a fibration sequence

$$B\Gamma_{\infty,n} \rightarrow B\Gamma_\infty^{(n)} \xrightarrow{\theta_n} B(S_n \wr GL^+(2, \mathbb{R})) \simeq B(S_n \wr S^1).$$

By forgetting marked points, one has a forgetful map $\varphi : B\Gamma_\infty^{(n)} \rightarrow B\Gamma_\infty$ so that the composite, $B\Gamma_{\infty,n} \rightarrow B\Gamma_\infty^{(n)} \xrightarrow{\varphi} B\Gamma_\infty$ is a homology isomorphism by Harer's stability theorem [2]. Therefore, φ induces a homology equivalence between this fibration and the trivial fibration,

$$\varphi : B\Gamma_\infty^{(n)} \xrightarrow{\cong_{H*}} B\Gamma_\infty \times B(S_n \wr S^1).$$

Upon applying Quillen's plus construction, this homology equivalence becomes a homotopy equivalence, and Theorem 11 follows.

Now the classifying space $B(S_n \wr S^1) \simeq ES_k \times_{S_k} (BS^1)^k = ES_k \times_{S_k} (\mathbb{CP}^\infty)^k$, and the classical result of Barratt-Quillen-Priddy is that when one takes the colimit over k , there is a homology equivalence

$$\mathbb{Z} \times \varinjlim_{k \rightarrow \infty} ES_k \times_{S_k} (\mathbb{CP}^\infty)^k \xrightarrow{\cong_{H*}} Q(\mathbb{CP}_+^\infty).$$

Passing to the Quillen plus construction we therefore have the following:

Corollary 12. *There is a homotopy equivalence*

$$\mathbb{Z} \times (B\Gamma_{\infty,n}^{(\infty)})^+ \simeq (B\Gamma_\infty)^+ \times Q(\mathbb{CP}_+^\infty).$$

Now recall Madsen and Weiss's theorem, proving a generalization of Mumford's conjecture on the cohomology of the stable mapping class group.

Theorem 13. *There is a homotopy equivalence*

$$\alpha : \mathbb{Z} \times (B\Gamma_\infty)^+ \xrightarrow{\simeq} \Omega^\infty(\mathbb{CP}_-^\infty).$$

of infinite loop spaces.

Combining Theorem 6, Corollary 12, and Theorem 13, we obtain Theorem 7.

3 Using more D -branes

Observe that taken together, Theorems 6 and 7 of the last section imply the main Theorem 1 in the case that the set of D -branes $\mathcal{B}$ consists of only one element. In this section we consider the case when we have a finite set of D -branes $\mathcal{B}$, and we describe the necessary modifications of the arguments in the last section needed to prove Theorem 1 in this general setting.

In the presence of a set $\mathcal{B}$ of D -branes, we define our general objects to be disjoint unions of circles and intervals, but now each interval has its boundary components labeled by D -branes. That is, we have the following definition.

Definition 7. *Given a finite set $\mathcal{B}$, a $\mathcal{B}$ -general object is a sequence $\vec{n} = (n_1, \dots, n_k)$ where each $n_i \in \{0, 1\}$, together with a labeling function of the boundary of the intervals of $C_{\vec{n}}$,*

$$\beta : \partial C_{\vec{n}} \rightarrow \mathcal{B}.$$

We similarly define a $\mathcal{B}$ -general morphism to be an open closed cobordism that respects the labeling functions.

Definition 8. *A $\mathcal{B}$ -general morphism from $(\vec{n}, \beta_1)$ to $(\vec{m}, \beta_2)$ is a general morphism Σ from $\vec{n}$ to $\vec{m}$ (in the sense of Definition 2), together with a labeling function of its free boundary components,*

$$\beta_\Sigma : \partial_f \Sigma \rightarrow \mathcal{B}$$

(where β_Σ is constant on each connected component) such that the free boundary $\partial_f \Sigma$ is a labeled cobordism from $\partial C_{\vec{n}} = \partial(\partial^- \Sigma)$ to $\partial C_{\vec{m}} = \partial(\partial^+ \Sigma)$. By “labeled cobordism” we mean that the labeling functions are compatible. That is,

$$\beta_1 \sqcup \beta_2 : \partial C_{\vec{n}} \sqcup \partial C_{\vec{m}} \rightarrow \mathcal{B}$$

is equal to the restriction of β_Σ to the boundary,

$$\beta_\Sigma : \partial(\partial_f \Sigma) \rightarrow \mathcal{B}.$$

As we did in section 1, we need to stratify the objects further by considering “open boundary permutations.” Namely, we observe that if (Σ, β_Σ) is a $\mathcal{B}$ -general morphism between $(\vec{n}, \beta)$ and (0) , (i.e an open-closed, $\mathcal{B}$ -labeled cobordism from $C_{\vec{n}}$ to (0)) then, as in definition 4 above, Σ defines an open boundary permutation $\sigma_\Sigma \in S_{\alpha(\vec{n})}$.

This allows us to define the open-closed string 2-categories $(\mathcal{S}_D^{\text{oc}})_{\mathcal{B}}$, $(\mathcal{S}_\Gamma^{\text{oc}})_{\mathcal{B}}$, and ultimately $\mathcal{S}_{\mathcal{B}}^{\text{oc}}$ just like in section one, but the objects will now be $\mathcal{B}$ -general objects together with open-boundary permutations, $(\vec{n}, \beta, \sigma)$. The 1-morphisms from $(\vec{n}, \beta, \sigma)$ to $(\vec{m}, \beta_2, \tau)$ are $\mathcal{B}$ -general morphisms (Σ, β_Σ) from $(\vec{n}, \beta)$ to $(\vec{m}, \beta_2)$ that pull back the open boundary permutations, $\Sigma^*(\sigma) = \tau$. The 2-morphisms $(\Sigma, \beta_\Sigma) \rightarrow (\Sigma', \beta_{\Sigma'})$ are the isomorphisms $\varphi : \Sigma \rightarrow \Sigma'$ of open-closed cobordisms as defined in Definition 3, with the additional requirement that they preserve the labeling functions:

$$\beta_{\Sigma'} \circ \varphi = \beta_\Sigma : \partial_f \Sigma \rightarrow \mathcal{B}.$$

The proof of Theorem 3 then gives the following generalization.

Theorem 14. *$\mathcal{S}_{\mathcal{B}}^{\text{oc}}$ is a strict symmetric monoidal 2-category.*

In the presence of the set $\mathcal{B}$ of D -branes, Theorem 6 needs the following modifications. Notice first of all that the windows in a $\mathcal{B}$ -labeled open-closed cobordism (Σ, β_Σ) (a 1-morphism in $\mathcal{S}_{\mathcal{B}}^{\text{oc}}$) are connected components of the free boundary, and therefore are equipped with labels in $\mathcal{B}$:

$$\beta_\Sigma : W \rightarrow \mathcal{B} \tag{2}$$

where $W \subset \pi_0(\partial_f(\Sigma))$ is the set of free boundary components which are windows. The set W is partitioned as

$$W = \sqcup_{b \in \mathcal{B}} W_b$$

where $W_b = \beta_\Sigma^{-1}(b)$. Let $w_b \in \mathbb{Z}^+$ be the cardinality of the set W_b . The analogue of Lemma 8 above is the following:

Lemma 15. *Let $\mathcal{BS}_{\mathcal{B}}^{\text{oc}}((\vec{m}, \beta, \tau), (0))$ be the classifying space of the category of morphisms from the object $(\vec{m}, \beta, \tau)$ in $\mathcal{S}^{\text{oc}}$ to the object (0) . Then*

$$\mathcal{BS}_{\mathcal{B}}^{\text{oc}}((\vec{m}, \beta, \tau), (0)) \simeq \coprod_{w=\{w_b\} \in (\mathbb{Z}^+)^{|\mathcal{B}|}} \coprod_{g \geq 0} B\Gamma_{g, c(\vec{m}, \beta, \tau)}^w.$$

Here the number $c(\vec{m}, \beta, \tau)$ is defined completely analogously to the description of $c(\vec{m}, \tau)$ prior to the statement of Lemma 8. The group $\Gamma_{g, c(\vec{m}, \beta, \tau)}^w$ is the mapping class group of isotopy classes of diffeomorphisms of a surface of genus g , with $c(\vec{m}, \beta, \tau)$ boundary components, and $w = \sum_{b \in \mathcal{B}} w_b$ marked points, where w_b of these marked points are labeled by the element b . These diffeomorphisms can permute these marked points, but must preserve the labelings.

One then defines a contravariant functor $(\mathcal{B}\mathcal{S}_{\mathcal{B}}^{\text{oc}})_{\infty}$ from $\mathcal{B}\mathcal{S}_{\mathcal{B}}^{\text{oc}}$ to spaces like what was done in Section 2. However in the presence of the set of D -branes $\mathcal{B}$ we define $\mathcal{B}(\mathcal{S}_{\mathcal{B}}^{\text{oc}})_{\infty}$ to be the homotopy colimit of the map

$$t_{\mathcal{B}} : \mathcal{B}\mathcal{S}^{\text{oc}}((\vec{m}, \beta, \tau), (0)) \rightarrow \mathcal{B}\mathcal{S}^{\text{oc}}((\vec{m}, \beta, \tau), (0))$$

defined by gluing on the surface $(T_{\mathcal{B}}, \beta_{T_{\mathcal{B}}})$ defined to be a morphism in $\mathcal{S}_{\mathcal{B}}^{\text{oc}}((0), (0))$ of genus one, with exactly one window W_b labeled by each element $b \in \mathcal{B}$. See Figure 5.




Figure 5: $T_{\mathcal{B}}$. The genus is one, and there is one window labeled by each $b \in \mathcal{B}$

The remaining part of the proof of Theorem 6 above goes through to prove the following analogue:

Theorem 16. *For each n there is a homology equivalence*

$$\mathbb{Z} \times \mathbb{Z}^{|\mathcal{B}|} \times B\Gamma_{\infty, n}^{\mathcal{B}} \xrightarrow{\cong_{H*}} \Omega|\mathcal{S}_{\mathcal{B}}^{\text{oc}}|.$$

where $B\Gamma_{\infty, n}^{\mathcal{B}}$ is defined as in the statement of Theorem 1.

Now just like Theorem 7 above is a consequence of Bødigheimer and Tillmann's splitting of the classifying spaces of stable mapping class groups (Theorem 3.1 of [1]) and Madsen and Weiss's theorem (Theorem 13 above) the following analogue also follows from these theorems.

Theorem 17. *For each n there is a homotopy equivalence*

$$\mathbb{Z} \times \mathbb{Z}^{|\mathcal{B}|} \times (B\Gamma_{\infty, n}^{\mathcal{B}})^+ \simeq \Omega^{\infty}(\mathbb{CP}_{-1}^{\infty}) \times \prod_{b \in \mathcal{B}} Q(\mathbb{CP}_{+}^{\infty})$$

Finally we notice that Theorems 16 and 17 together are equivalent to the statement of Theorem 1.

References

- [1] C.F. Bødigheimer and U. Tillmann, *Stripping and splitting decorated mapping class groups*, Progress in Math. **(78)**, Birkhauser (2001), 47 - 57.
- [2] J.L. Harer, *Stability of the homology of the mapping class groups of orientable surfaces* Ann. Math. **121**, (1985), 215-249.
- [3] I. Madsen and M. Weiss, *The stable moduli space of Riemann surfaces: Mumford's conjecture*, preprint arXiv:math.AT/0212321 (2004)

- [4] D. McDuff and G. Segal, *Homology fibrations and the group completion theorem*, Invent. Math. **31**, (1976), 279-284.
- [5] G. Segal, *Categories and cohomology theories*, Topology **13**, (1974), 293-312.
- [6] G. Segal, *Topological structures in string theory*, Phil. Trans. R. Soc. Lond. A **359**, (2001), 1389-1398.
- [7] G. Segal, *The definition of conformal field theory*, Topology, Geometry, and Quantum Field theory, Proc. of the 2002 Oxford symposium in honour of the 60th birthday of Graeme Segal, LMS Lecture Notes Series **308**, Part II, (2004).
- [8] U. Tillmann, *On the homotopy of the stable mapping class group*, Invent. Math. **130**, no. 2 (1997), 257-275.
- [9] U. Tillmann, *Discrete models for the category of Riemann surfaces*, Math. Proc. Cam. Phil. Soc. **121**, (1997), 39 - 49.